

## Nutritional and Nutraceutical Properties of Pumpkin

Nikhil Thakur, Neeraj Gupta and Damanpreet Kaur  
Division of Food Science and Technology, SKUAST Jammu, 180009

ARTICLE ID: 26

### Introduction:

Originally from Central to South America, the trailing plant *Cucurbita*, also known as "pumpkin," is grown anywhere from sea level to high altitudes (Yadav *et al.* 2010). It is considered an indigenous vegetable in many other nations, including Africa. India, China, Mexico, and the United States are the major producers of pumpkin on the international market. Pumpkin is a member of the Cucurbitaceae family and the genus *Cucurbita*. There are five different varieties of pumpkin, but the three most prevalent are *Cucurbita pepo*, *Cucurbita maxima*, and *Cucurbita moschata* (Saeleaw and Schleining, 2011). There are several dietary benefits and food uses for pumpkin. Due to its high concentration of  $\beta$ -carotene and moderate number of carbohydrates, as well as its vitamins B6, K, thiamine, and riboflavin and minerals potassium, phosphorus, magnesium, iron, and selenium, it can be used as a food additive in the production of juice, pomade, pickles, and dried goods (Rakcejeva *et al.* 2011). Because they are so rich in micro and macro nutrients like phosphorous, calcium, magnesium, manganese, copper, and zinc, pumpkin seeds are frequently utilised as dietary supplements. The medical benefits of pumpkin include anti-inflammatory, antioxidant, anticarcinogenic, anti-angiogenesis, anti-lipogenic, glucose-lowering, and anti-cancer characteristics. The presence of many phytochemicals, including tocopherol ( $\alpha$  and  $\gamma$ -tocopherol), carotenoid (lutein, zeaxanthin,  $\beta$ -carotene), triterpenes, and secondary metabolites such  $\beta$ -sitosterol, dehydrodiconiferyl alcohol, and tetra saccharide glycerol glycolipid, contributes to the abundance of health advantages (Caili *et al.* 2006; Kim *et al.* 2012; Wang *et al.* 2012; Isutsa and Mallowa, 2013).

### Pumpkin nutritional components:

The fruit of the *Cucurbita* genus is a superior source of nutrients. Pumpkin has high levels for pro-vitamin A, carotenoids, that give its distinctive colours due to the presence of substances like lutein (a bright yellow colour) and beta-carotene (orange color). The presence of  $\beta$ -carotene, lutein, lycopene and cryptoxanthin in trace levels is the exception (Zhou *et al.*

2007).  $\beta$ -Carotene is transformed to vitamin A within the body, which is essential for vision, growth, and embryonic development and a deficit may result in vision and infant mortality. By scavenging free radicals as it quenches singlet oxygen and interacts with the radical itself, carotenoid works as an antioxidant and lowers the chance of developing certain diseases like cancer and cardiovascular diseases (CVDs). Mostly made up of pulp and seeds, pumpkin fruit is rich in both necessary and non-essential amino acid residues, which are necessary for both bodily and mental functions (Takahashi *et al.* 2011). Lysine is the amino acid that is present in the highest concentrations among the essential amino acids, while glutamic acid is the case for the non-essential ones. On a dry weight basis, the pumpkin seed has a crude protein content of 39.25 g/100 g, a crude oil content of 27.83 g/100 g, an ash content of 4.59 g/100 g, a crude fibre content of 16.84 g/100 g, an energy content of 2,401 kcal per 100 g, and concentrations of potassium, chromium, sodium, magnesium, aluminium, barium, calcium, zinc, and iron of (Kim *et al.* 2012). On a dry weight basis, the pulp contains low amounts of fat (2.3 g/100 g), carbs (66 g/100 g), protein (3 g/100 g) and carotenes (171.9 mg/g) (Martha and Gutierrez, 2016). Pumpkin pulp contains 2-10 mg/100 g of vitamin C and 9-10 mg of vitamin E per 100 g.

#### **Nutraceutical properties:**

As an antirheumatic (a substance that reduces rheumatic disease symptoms), demulcent (a substance that shapes a soothing film over mucous membranes), diuretic (a substance that increases urine production), nervine (a calming tonic with pumpkin 387 properties that soothes frazzled nerves), and taenifuge (a medication to remove tapeworms from the body), *Cucurbita maxima* has a variety of beneficial effects (Bhat *et al.* 2015). Additionally, the fibres in pumpkin help to bind the extra acids that the digestive system produces, which helps to buffer the pH of the stomach. It helps regulate blood sugar levels while treating gastrointestinal parasites, supporting benign prostatic hyperplasia (BPH), supporting urinary dysfunctions, and dysuria (Vergara *et al.* 2006).

#### **Anti-oxidant activity:**

The assessment of antioxidant activity in Cucurbitaceae seed protein often includes a variety of in-vitro tests, including measures of metal chelating activity, reducing power, and radical scavenging activity (Ozuna and Leon Galvan, 2017). The isomeric combination of  $\beta$

or  $\delta$ -tocopherol is present in pumpkin seed oil (Matus *et al.* 1993). Tocopherols are a non-glycoside molecule found in vegetable oils that function as natural antioxidants.

**Hypotensive activity:**

Low blood pressure, particularly inside the arteries of a systemic circulation, is the main cause of hypotension. The findings demonstrated a preventive impact against pathological cardiac and aortic changes as well as lower chances of heart attacks due to the high magnesium content. Diastolic blood pressure significantly dropping in recent clinical research in postmenopausal women (Yoshinari *et al.* 2015).

**Antidiabetic activity:**

Due to insulin's non-secretion or resistance, diabetes mellitus is indeed a long-lasting metabolic condition that manifests as insulin shock (Makni *et al.* 2011). The pumpkin seed includes tocopherol isomers, which are useful in reducing diabetes since they have some antioxidant action. For intake in diabetic conditions, pumpkin fruit is employed (Xia and Wang, 2007).

**Anticancer activity:**

According to research, both antioxidants and polyphenolic substances in diet can regulate abnormal inflammatory signals and signalling pathways linked to cancer stem cells. The extract from cucurbita pepo may one day be used as a novel chemotherapy drug to treat cancer and tumour growth. Carotenoids from a diet high in pumpkin seeds have also been linked to a reduced risk of prostate cancer in addition to lowering the chances of gastric, breast, lung, and colon cancer. By eliminating a variety of free radicals produced in the body throughout metabolism, including as superoxide anions, hydroxyl radicals, and other reactive oxygen species, pumpkin polysaccharides have beneficial anticancer benefits (Chen and Huang, 2019).

**Antimicrobial activity:**

Pumpkin is significant for clinical microbiology and has medicinal uses. For pharmacological and microbiological studies, Cucurbita maxima leaves, fruit, and flower extract is used. Pseudomonas aeruginosa, Candida albicans, Acinetobacter baumannii, Enterococcus faecalis, Klebsiella pneumonia, Escherichia coli, and Staphylococcus aureus are all susceptible to the antibacterial properties of pumpkin oil-soluble components. The examined microbial isolates were resistant to three distinct proteins derived from pumpkin

rinds, seeds, and pulp. Aqueous extract against the fungus *Aspergillus flavus* and the toxicity of aflatoxin B1 (AFB1) Researchers looked into the nutritional benefits of pumpkin 391 infection-related lung histolo morphological damages in rats (Saddiq *et al.* 2019).

#### References:

- Bhat, A., Satpathy, G. and Gupta, R. K. (2015), Evaluation of nutraceutical properties of *amaranthus hypochondriacus* L. grains and formulation of value added cookies, *Journal of Pharmacognosy and Phytochemistry*, **3**(5): 51-54.
- Caili, F. U., Huan, S. and Quanhong, L. I. (2006), A review on pharmacological activities and utilization technologies of pumpkin, *Plant Foods for Human Nutrition*, **61**(2): 70-77.
- Chen, L. and Huang, G. (2019), Antioxidant activities of sulfated pumpkin polysaccharides, *International Journal of Biological Macromolecules*, **126**: 743-746.
- Isutsa, D. K. and Mallowa, S. O. (2013), Increasing leaf harvest intensity enhances edible leaf vegetable yields and decreases mature fruit yields in multi-purpose pumpkin, *Journal of Agricultural and Biological Science*, **8**: 610-615.
- Kim, M. Y., Kim, E. J., Kim, Y. N., Choi, C. and Lee, B.H. (2012), Comparison of the chemical compositions and nutritive values of various pumpkin (*Cucurbitaceae*) species and parts, *Nutrition Research and Practice*, **6**(1): 21-27.
- Makni, M., Fetoui, H., Gargouri, N. K., Garoui, E. M. and Zeghal, N. (2011), Antidiabetic effect of flax and pumpkin seed mixture powder: effect on hyperlipidemia and antioxidant status in alloxan diabetic rats, *Journal of Diabetes and Its Complications*, **25**(5): 339-345.
- Martha, R. and Gutierrez, P. (2016), Medicinal chemistry review of *Cucurbita pepo* (pumpkin) its phytochemistry and pharmacology, *International Journal of Pharmacology and Phytochemistry Research*, **9**(9): 1190-1194.
- Matus, Z., Molnar, P., Szabo, L. J. (1993), Main carotenoids in pressed seeds (*Cucurbita semen*) of foil pumpkin (*Cucurbita pepo* convar. *Pepo*. Var. *styriaca*), *Acta Pharmaceutica Hungarica*, **63**(5): 247-256.
- Ozuna, C. and Leon-Galvan, M. F. (2017), Cucurbitaceae seed protein hydrolysates as a potential source of bioactive peptides with functional properties, *BioMedical Research International*, **2017**: 2121878.

- Rakcejeva, T., Galoburda, R., Cude, L. and Strautniece, E. (2011), Use of dried pumpkins in wheat bread production, *Procedia Food Science*, **1**: 441-447.
- Saddiq, N., Ali, A. and Awedh, M. H. (2019), Pumpkin (*Cucurbita moschata*) against *Aspergillus flavus* and aflatoxin B1 induced lung cyto-morphological damage in rats, *Pakistan Journal of Pharmaceutical Sciences*, **32**(2): 575-579.
- Saeleaw, M. and Schleining, G. (2011), Composition, physicochemical and morphological characterization of pumpkin flour, Proceeding of the 11th *International Congress on Engineering and Food*, pp. 10-13.
- Takahashi, T., Toda, E., Singh, R. B., De Meester, F., Wilczynska, A., Wilson, D. and Juneja, L. R. (2011), Essential and nonessential amino acids in relation to glutamate, *The Open Nutraceuticals Journal*, **4**(1): 205-212.
- Vergara-Valencia, N., Granados-Perez, E., Agama-Acevedo, E., Tovar, J., Ruales, J. and Bello-Perez, L. A. (2006), Fiber concentration from mango fruit characterization associated antioxidant capacity and application as a bakery product ingredient, *Food Science and Technology*, **40**(4): 722-729.
- Wang, S. Y., Huang, W. C., Liu, C. C., Wang, M. F., Ho, C. S., Huang, W. P., Hou, C. C., Chuang, H. L. and Huang, C. C. (2012), Pumpkin (*Cucurbita moschata*) fruit extract improves physical fatigue and exercise performance in mice”, *Molecules*, **17**(10): 11864-11876.
- Xia, T. and Wang, Q. (2007), Hypoglycaemic role of *Cucurbita ficifolia* (Cucurbitaceae) fruit extract in streptozotocin-induced diabetic rats, *Journal of the Science of Food and Agriculture*, **87**(9): 1753-1757.
- Yadav, M., Jain, S., Tomar, R., Prasad, G. B. K. S. and Yadav, H. 2010. Medicinal and biological potential of pumpkin: an updated review. *Nutrition research reviews*, **23**(2): 184-190.
- Yoshinari, O., Udani, J., Moriyama, H., Shiojima, Y. and Chien, X. (2015), The efficacy and safety of a proprietary onion-pumpkin extract (OPtain120) on blood pressure: an open-label study, *Functional Foods in Health and Disease*, **5**(6): 224-242.
- Zhou, T., Kong, Q., Huang, J., Dai, R. and Li, Q. (2007), Characterization of nutritional components and utilization of pumpkin, *Food*, **1**: 313-321.